



TITLE:

Local structure and dynamics in colloidal fluids, clusters and gels(Poster session 2, New Frontiers in Colloidal Physics : A Bridge between Micro- and Macroscopic Concepts in Soft Matter)

AUTHOR(S):

Ohtsuka, Takehiro; Royall, Paddy; Tanaka, Hajime

CITATION:

Ohtsuka, Takehiro ...[et al]. Local structure and dynamics in colloidal fluids, clusters and gels(Poster session 2, New Frontiers in Colloidal Physics : A Bridge between Micro- and Macroscopic Concepts in Soft Matter). 物性研究 2007, 89(1): 158-158

ISSUE DATE:

2007-10-20

URL:

<http://hdl.handle.net/2433/110891>

RIGHT:

Local structure and dynamics in colloidal fluids, clusters and gels

Inst. of Industrial Science, The Univ. of Tokyo Takehiro Ohtsuka, Paddy Royall ^{1 2}
and Hajime Tanaka

Gels in soft matter systems often exhibit hallmarks of arrested phase separation. We consider a colloid-polymer mixture which is quenched by increasing the polymer concentration, from a fluid to a ‘cluster fluid’ and then a non-ergodic gel. Using confocal microscopy, we study both the static structure and dynamics in 3D, presenting the first time-resolved 3D coordinate tracking of a colloidal fluid. While significant changes in structure are observed upon the formation of a network, they are accompanied by a limited dynamic slowing.

Furthermore, deeper quenching, upon which there is a significant dynamic slowing, is accompanied by more subtle changes in structure. We find evidence of five-fold symmetry, which may inhibit access to global potential energy minimum, fluid-crystal coexistence, leading to gelation. We find strong spatial correlations between particle motion at all state points, including the non-ergodic gel. In particular, the connection between coordination number and (local) displacement suggests mobility enhancement on the gel surface, hinting at ‘surface melting’.

¹E-mail: paddy@iis.u-tokyo.ac.jp

²Also at the University of Bristol, UK.